

Comparative Phytochemical Screening and Pharmacognostic Study on Stem Bark and Root bark of Endangered Medicinal Plant *Oroxylum indicum* (L.) Vent

Virendra Rajak^{1*}, Vandana Pathak², Manoj Tripathi³

^{1,2}Mahatma Gandhi Chitrakoot Gramodaya, Vishwavidyala, Chitrakoot, Satna, India

³Arogyadham (J.R.D. Tata Foundation for Research in Ayurveda & Yoga Sciences),
Deendayal Research Institute, Chitrakoot, Dist Satna, India

*Correspondence Author: virendrarajak007@gmail.com

Available online at: www.isroset.org

Received: 02/May/2018, Revised: 14/May/2018, Accepted: 12/Jun/2018, Online: 30/Jun/ 2018

Abstract- Medicinal plants are an important component of natural resources and they are used as a source of food, medicine, fibres, tannin, gum and allied uses. *Oroxylum indicum* (L.) Vent. (*Bignoniaceae*) also known as 'Sonapatha' is an important endangered medicinal plants of indigenous medical system for over hundreds of years. It has been used as a single drug and as a component for preparation of ayurvedic compound preparations viz. Dashmularistha and Chyavanprash etc. Plants are used in treating in various human ailments like jaundice, dysentery, asthma, allergic disease, urticaria, sore throat, measles, laryngitis, hoarseness and diarrhea. The present paper provides a detailed account of the phytochemical evaluation and pharmacognostic studies of Sonapatha stem bark and root bark. The study includes macroscopy, microscopy, and powder microscopy studies, fluorescence study, physicochemical parameters, preliminary phytochemical screening High Performance Thin Layer Chromatography fingerprint profile. Established parameters can be used as standards for identification and quality control of the plants in compound formulations and also preparation of a monograph of the plant.

Keywords- *Oroxylum indicum*, , Phytochemical screening, Physico-chemical studies, HPTLC fingerprints profile

I. INTRODUCTION

Medicinal plants play an important role in the prevention and treatment of various human ailments since time immemorial (Samanta et al., 1999). In Indian system of medicines a large number of herbal and herbo-mineral drugs have been used to cure various types of diseases. Ayurveda is one of the traditional systems of medicine practiced in India back to 6000 BC. (Gupta et al., 2006). *Oroxylum indicum* (L.) Vent. (*Bignoniaceae*) also known as 'Sonapatha' is an important endangered medicinal plants. It is native to the Indian subcontinent, in the Himalayan foothills with a part extending to Bhutan and southern China, in Indo-China and the Malaysia ecozone. It is diversely available in the forest of National Park in Assam, India, reported from Sri Lanka (Ceylon) (Kirtikar and Basu, 2001). It is large medium size tree 12-15 meter (40 feet) height. Stem bark colour is off brown. Leaves are 2.5 to 5 inch long, broad, leaflets are 4.5 inch long and 3.5 to 4.5 inch broad having sharp edges. Flowers colour is purple and stalk is 0.5-1.5 feet long. Fruits long curve downward and resemble the wings of a large bird or dangling sickles or swords in the night and 1 to 3.5 feet long, 2.5 to 4.5 inch broad. Seeds are flat 2.5-3.5 inch long and 1.5-2-5 inch in

wide. The flowers are born in rainy season and fruiting session in December to March (Dalal and Rai 2004). *Oroxylum indicum* used to various Ayurvedic preparations viz. Chyavanaprasha Awaleha, Amartarista, Dasamula, Dhanawantara Ghrita, Dantyadyarista, Brahma Rasayana, Narayana Taila and mentat (mental drug) etc. (Balkrishna, 2005; Kumar et al., 2009; Anonymous, 1998). Sonapatha different parts are used as tannins, dyestuffs and wood (Laupattarakasem et al., 2003; Gupta et al., 2008). *Oroxylum indicum* stem bark decoction is used to cure cancer (Mao, 2002). In various tribes of India, bark and seeds of the plant are used in fever, stomach disorders, pneumonia and respiratory troubles (Panghal et al., 2010; Patil et al., 2008, Raut et al., 2009). The purpose of this study there are no reports of systematic pharmacognostic investigation of stem bark and stem bark of this plant. So task was undertaken to scientific evaluation of endangered medicinal plant Sonapatha stem bark and root bark. The study includes macro and microscopic characters, powder microscopic characteristics, High Performance Thin Layer Chromatography fingerprints, preliminary phytochemical screening and physicochemical parameters. The information generated by this particular study provides relevant pharmacognostical and phytochemical data needed for

proper identification and authentication of Sonapatha stem bark and root bark.

II. MATERIAL AND METHODS

Collection and processing of plant material:

The fresh plant stem bark and root bark of *Oroxylum indicum* was collected from the, Sati-anusuiya forest of Chitrakoot, Satna (M.P.) in the month of March. Samples were authenticated (Jain 1991, Kirtikar and Basu 1935, Verma *et al.*1993) by Botanist Dr. Manoj Tripathi of Ayurveda Sadan (Research Laboratory), Deendayal Research Institute Chitrakoot. The voucher specimen (AD/AS/124/2016) prepared as per standard procedure (Jain and Rao 1977) and maintained in the herbarium of faculty of Science and Environment, Mahatma Gandhi Chitrakoot Gramodaya University, Chitrakoot, Satna (M.P.) for further reference. Fresh material was used for anatomical studies whereas shade dried material was powdered in electric grinder for physico-chemical study, phytochemical investigation and development of High Performance Thin Layer Chromatography fingerprint profile.

Macroscopic study:

Macroscopic or organoleptic characters like appearance, colour, odour and taste were evaluated.

Powder microscopic study:

The dried stem bark and root bark were subjected to powdered and completely passes through 355 μm IS Sieve (old sieve number 44) and not less than 50% pass on through 180 μm IS Sieve (old sieve number 85). About 2 g of powder washed thoroughly with potable water, pour out the water without loss of material. Mounted a small portion in glycerine, warmed a few mg with chloral hydrate solution, wash and mounted in glycerin. Treat a few mg with iodine solution and mount in glycerine, about 1 g of powder warmed over water bath with Chloral hydrate solution till brown fumes appear, cool and wash with water thoroughly and mount a small portion in glycerin and seen under microscope at 40X x 10X magnification of the Trinocular Research Microscope (Evans 2003 and Kokate 2006).

Physico-chemical parameters:

Physico-chemical parameters such as moisture content (loss on drying at 105⁰C), water soluble extractive value, alcohol soluble extractive value, total ash value, acid insoluble ash value and water soluble ash were calculated (Anonymous 2000 and Anonymous 2007).

Preliminary phytochemical studies:

Preliminary phytochemical tests were carried out on ethanolic and water extract for the presence/absence of phyto-constituents like alkaloids, flavanoids, steroids, terpenoids, tannins, resins, carbohydrates, proteins and saponins (Tripathi and Sikarwar 2015, Tripathi *et al.* 2015).

High Performance Thin Layer Chromatography (HPTLC):

For High performance thin layer chromatography, the powdered 5 gm of each sample was extracted with 100 ml of ethanol overnight, filtered and concentrated. It was applied by spotting extracted sample on pre-coated silica-gel aluminium plate 60 F₂₅₄ (5x10 cm with 0.2 mm layer thickness Merk Germany) using Camag Linomat -5 sample applicator and a 100 μl Hamilton syringe. The samples, in the form of bands of length 6 mm, were spotted 15 mm from the bottom, 15 mm from left margin the plate and 10 mm part. Plates were developed using mobile phase consisting of *Toluene: Ethyl acetate* (7:3 v/v). Linear ascending development was carried out in 10x10cm twin through glass chamber equilibrated with mobile phase. The optimized chamber saturation time for mobile phase was 30 min. at room temperature. The length of chromatogram run was 8 cm. 20 ml of the mobile phase. Subsequent to the development, Thin Layer Chromatography plate was dried with the help of Hot Air Oven. The peak area for samples and standard were recorded with Camera photo documentation system Camag Reprostar 3. Visualization of spot was made before and after derivatization (with 5% Methanolic - sulphuric acid reagent) at 254nm, 366nm and day light with Win cat software and R_f values noted (Ansari 2006, Choudhary *et al.* 2014).

III. RESULTS AND DISCUSSION

Macroscopic, microscopic and powder microscopic study:

Sonapatha stem bark colour is off brown, taste acrid, bitter, odour astringent. Detailed Transverse Section of Sonapatha stem bark shows mu8-12 layered cork cells, cortex wide consisting thick walled rectangular parenchymatous cells some are embeded with prismatic crystals of calcium oxalate, wide lumen, round to hexagonal stone cells present in the cortex. medullary rays present. Stem bark powder examined under microscope shows cork cells in sectional view, thick walled cork cells in surface view, parenchymatous cells, thick walled with wide lumen tetragonal to hexagonal group of stone cells, hick walled long fibres, prismatic crystals of calcium oxalate, longitudinally cut medullary rays and transversely cut medullary rays with parenchymatous cells.

Sonapatha dried root pieces are cylindrical stout, woody and light, 5 to 9 cm length, 4 to 6 cm in diameter, surface smooth, or often uneven, marked with few distantly placed lenticels, many vertically placed narrow streaks or transverse cracks at places often faintly fissured, shrunken and longitudinally wrinkled, at places the outermost thin bark gets separated exposing the inner green coloured bark adherent to the inner buff coloured porous wood. Odour characteristics, taste slightly bitter, mucilaginous. Detailed Transverse Section of Sonapatha root bark shows mu14-18

layered cork cells, cortex wide consisting thick walled rectangular parenchymatous cells some are embedded with prismatic crystals of calcium oxalate, wide lumen, cylindrical to hexagonal stone cells present in the cortex. medullary rays present. Root bark powder examined under microscope shows cork cells in surface view, thick walled cork cells in sectional view, cortical parenchymatous cells, pitted parenchymatous cells, thin walled with wide lumen group of stone cells, reticulate and simple pitted thickenings, thick walled phloem fibres, prismatic crystals of calcium oxalate and acicular crystals




Figure-1: Sonapatha plant

Physico-chemical analysis:

The physico-chemical parameters such as extractive values are useful for the determination of exhausted or adulterated drug; ash values of the drug gave an idea of the earthy matter or the inorganic composition and other impurities present along with the drug. Physico-chemical results of the drug are given in (Table 1).

Table 1: Physico-chemical analysis of the Sonapatha stem bark and root bark

S. No.	Parameters	Results stem bark	Root bark
1	Foreign matter	1.9%	1.6%
2	Loss on drying at 1050C	5.24%	6.35%
3	Ethanol-soluble extractive	14.50%	23.30%
4	Water-soluble extractive	21.15%	33.15%
5	Total ash value	11.4%	10.50%
6	Acid-insoluble ash value	2.4%	2.15%

Preliminary phyto-chemical investigation:

Qualitative phyto-constituents were screened in the extracts taken in water and ethyl alcohol. Phyto-chemical results of the drug are given in (Table 2).

Table 2: Phytochemical analysis of Sonapatha stem bark and root bark

S. No.	Name of phyto-constituents	Result Stem bark	Root bark
1	Alkaloids	Present	Present
2	Carbohydrates	Present	Present
3	Protein	Present	Present
4	Saponin	Present	Absent
5	Steroids	Present	Present
6	Resin	Absent	Present
7	Tannin	Present	Absent
8	Flavonoid	Absent	Present

High performance thin layer chromatography:

High performance thin layer chromatography (HPTLC) study of the ethanolic extract two spots of the sample extracts applied in the Thin Layer Chromatography plate. Major spots R_f values with colour were recorded under, 254nm, 366nm, after derivatization 366nm. Chromatogram profile and R_f values are given (Figure.12 -14and Table -3).

The macroscopic, microscopic and powder microscopic distinguished characters have been established to identify Sonapatha stem bark and root bark. The pharmacognostic and physicochemical parameters can be used for checking the adulteration and purity of this drug. High performance thin layer chromatography finger print profile helps in identification of various phytochemical constituents present in the crude drug thereby substantiating and authenticating of crude drug. The High performance thin layer chromatography profile also helps to identify and isolate's important phyto-constituents. These finding could be helpful in identification and authentication.

Table-3: R_f values of HPTLC finger prints stem bark

R_f values	Test solution of Sonapatha stem bark	
	At 366 nm(before derivatization)	At 366nm (after derivatization)
R_{f1}	0.04(light pink)	0.06 (light pink)
R_{f2}	0.08(dark pink)	0.18(light brown)
R_{f3}	0.32(brown)	0.32(yellow)
R_{f4}	0.34(reddish brown)	0.34(sky blue)
R_{f5}	0.60(dark red)	0.60(yellow)

Table-4: R_f values of HPTLC finger prints root bark

R_f values	Test solution of Sonapatha root bark	
	At 366 nm(before derivatization)	At 366nm (after derivatization)
R_{f1}	0.14 (sky blue)	0.10 (whitish blue)

R _f 2	0.38 (florescence)	0.16(whitish blue)
R _f 3	0.58 (florescence)	0.38 (yellow)
R _f 4	0.74(florescence)	0.58 (yellow)
R _f 5	0.78(red)	0.74(yellow)




Figure2. HPTLC fingerprint profile At 366nm before derivatization stem bark



Figure-3. HPTLC fingerprint profile At 366nm after derivatization stem bark




Figure4. HPTLC fingerprint profile At 366nm before derivatization root bark




Figure-5. HPTLC fingerprint profile At 366nm after derivatization root bark

IV. CONCLUSION

Oroxylum indicum has numerous uses in traditional medicine to treat several ailments viz. in treating cancer, fever, stomach disorders, pneumonia and respiratory troubles etc. It is also used to various Ayurvedic preparations viz. Chyavanaprasa Awaleha, Amartarista, Dasamula, Dhanawantara Ghrita, Dantadyarista, Brahma Rasayana, Narayana Taila and mentat (mental drug) etc. Due to its wide therapeutic importance it is worthwhile to standardize it for use as drug. The present study reveals phytochemical screening and pharmacognostic study of drug *Sonapatha*, which would be of immense value in botanical identification and authentication of plant drug may help us in preventing its adulteration.

V. ACKNOWLEDGEMENT

The authors are grateful to the Prof. N. C. Gautam, Vice Chancellor, Mahatma Gandhi Chitrakoot Gramodaya, university Chitrakoot, Satna (M.P.) for providing necessary facilities.

REFERENCE

- [1]. Anonymous, The Ayurvedic Pharmacopoeia of India. Part-I, Volume-III, Government of India, Ministry of Health and Family Welfare, Deptt. of ISM & H., The Controller of Publications Civil Lines, Delhi, 2000.
- [2]. Anonymous. The Ayurvedic Pharmacopoeia of India, Government of India, Ministry of health and family welfare Department of Indian system of medicine and Homeopathy, New Delhi, India. 1998; 209-210.
- [3]. Anonymous, Protocol for Testing of Ayurvedic, Siddha & Unani Medicines. Government of India, Department of AYUSH, Ministry of Health & Family Welfare, Pharmacopoeial Laboratory for Indian Medicines Ghaziabad, 2007.
- [4]. Ansari SH, *Essentials of Pharmacognosy* 1st edition, Birla publications Pvt. Ltd, New Delhi, 2006.
- [5]. Balkrishna A. Ayurved jadi-booti rahasya. Divyaprakasan, Divya yog mandir trust, Haridwar, Uttarakhand. 2005; 132-136.
- [6]. Choudhary N, Siddiqui MB and Khatoon S, Pharmacognostic evaluation of *Tinospora cordifolia* Willd.) Miers and identification of biomarkers, *Indian J Tradit Knowle*, 2014 ; 13(3) 543-540.
- [7]. Dalal NV, Rai VR. In vitro propagation of *Oroxylum indicum* Vent. A medicinally important forest tree. *J For Res* 2004;9:61-65.
- [8]. Evans WC, Trease and Evans Pharmacognosy, 15th ed., (Saunders, London, 2003, 545-547.
- [9]. Gupta M, Mazumder UK, Gomathi P, Thamil SV (2006). Antiinflammatory evaluation of leaves of *Plumeria acuminata*, *BMC Complementary and alternative Medicine*, 6: 36.
- [10]. Gupta R.C., Sharma V., Sharma N., Kumar N., Singh B. In vitro antioxidant activity from leaves of *roxylum indicum* (L.)(Vent.)- A north Indian highly threatened and vulnerable medicinal plant. *J Pharm Res*. 2008; 1(1): 65-72.
- [11]. Jain SK, *Dictionary of Indian Folk Medicine and Ethnobotany*. Deep Publications, New Delhi, 1991.
- [12]. Jain SK and Rao RR, *A hand book of field and herbarium methods*, Today and tomorrow printers and publications New Delhi 1977.

- [13]. Kirtikar KR, Basu BD. Indian Medicinal Plants. Oriental Enterprises, Dehradun, Vol. 4 2001; 1105–1107.
- [14]. Kirtikar KR and Basu BD, *Indian Medicinal Plants*. Bishen Singh Mahendra Pal Singh, Dehra Dun 1935, Vols. 1(4) (2nd ed.).
- [15]. Kokate CK, Practical Pharmacognosy Is ted., Vallabh Prakashan, New Delhi 2006.
- [16]. Kumar V., Gogoi B.J., Meghvansi M.K., Singh L., Srivastava R.B., Deka D.C. Determining the antioxidant activity of certain medicinal plants of Sonitpur, (Assam), India using DPPH assay. *J Phyto.* 2009; 1(1): 49–56.
- [17]. Laupattarakasem P., Houghton P. J., Hoult J. R., Itharat A. An evaluation of the activity related to inflammation of four plants used in Thailand to treat arthritis. *J Ethnopharmacol.* 2003; 85(2–3): 207–215.
- [18]. Mao A, *Oroxylum indicum* Vent- A potential anticancer medicinal plant. *Indian Journal of Traditional Knowledge* 2002, 1(1): 17-21.
- [19]. Patil G.G., Mali P.Y., Bhadane V.V. Folk remedies used against respiratory disorders in Jalgaon district, Maharashtra. *Nat Prod Rad.* 2008; 7(4): 354-358.
- [20]. Panghal M., Arya V., Yadav S., Kumar S., Yadav J.P. Indigenous knowledge of medicinal plants used by Saperas community of Khetawas, Jhajjar district, Haryana, India. *J Ethnobi and Ethnomed.* 2010; 6: 4-11.
- [21]. Rout S.D., Panda T., Mishra N. Ethno-medicinal plants used to cure different diseases by tribals of Mayurbhanj District of North Orissa. *Ethno Medicine.* 2009; 3(1): 27-32.
- [22]. Samanta MK, Srinath, Pulok, KM, Suresh B. (1999). Herbal Drug Development- A modern Approach, *Ind. J. Pharma Edu.* 33(4): 202-204.
- [23]. Tripathi M and Sikarwar RLS, Pharmacognostic Studies on Plaksa (*Ficus virens* Ait.) Stem Bark, *Indian Journal of Natural Products and Resources* 2015; 6(1) 27-32.
- [24]. Tripathi M, Sikarwar RLS, Tiwari A and Dwivedi N., Pharmacognostical identification of ingredients in Laghulaiurna, *Indian Journal of Traditional Knowledge*, 2015; 14(4) 531-536.
- [25]. Verma DM, Balakrishnan NP and Dixit RD, *Flora of Madhya Pradesh Botanical Survey of India, Calcutta* 1993; Vol. (I).